

Radiation effects in  $\alpha$ -quartz

S/070/63/008/002/003/017  
E039/E435

quartz; d) the intensity of scattering in the irradiated quartz depends on the orientation of the crystal. The X-ray analysis shows that the third order symmetry  $C_3$  is changed to sixth order  $C_6$  by the irradiation and there is a significant change in the distribution of diffuse scattering. As a result of neutron irradiation, the structure of  $\alpha$ -quartz is thought to change in the following manner: 1) Initially, defects develop which lead to a weakening and breaking of the Si-O bond and hence to the possibility of rearrangement in the Si-O tetrahedrons. 2) At a definite stage of the exposure the  $\alpha$ -quartz becomes unstable and there is a transition to the more symmetrical high temperature modification. This remains stable at room temperature. 3) There is a complete loss of orientation in parts of the crystal. There are 4 figures.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet im.  
M.V.Lomonosova (Moscow State University imeni  
M.V.Lomonosov).

SUBMITTED: July 10, 1962  
Card 2/2

KOLONTSOVA, Ye.V.; MIKHAYLOVA, L.K.

Two-dimensional defects in plastically deformed crystals.  
Kristallografiia 8 no.6:900-905 N-D'63. (MIRA 17:2)

1. Moskovskiy gosudarstvennyy universitet imeni M.V. Lomonosova.

ZHDANOV, G.S.; ZUBOV, V.G.; KOLONTSOVA, Ye.V.; OSIPOVA, L.P.;  
TELEGINA, I.V.

Radiation effects in  $\alpha$ -quartz. Kristallografiia 8 no.2:207-212  
Mr-Apr '63. (MIRA 17:8)

1. Moskovskiy gosudarstvennyy universitet imeni Lomonosova.

ACCESSION NR: AP4024997

S/0070/64/009/002/0282/0284

AUTHORS: Kolontsova, Ye. V.; Telegina, I. V.

TITLE: Change in the structure of pentaerythrite when irradiated with x-rays

SOURCE: Kristallografiya, v. 9, no. 2, 1964, 282-284

TOPIC TAGS: pentaerythrite, x ray, pentaerythrite structure, irradiation, irradiate crystal, reflection intensity, point defect, dislocation, dislocation loop, fragmentation

ABSTRACT: It has been found that the intensity of reflected monochromatic rays from pentaerythrite does not remain constant but increases slightly at first and then declines. The "useful lifetime" of a crystal, when the intensity of reflection is still rather high and to some degree constant, varies from crystal to crystal, generally ranging from 50 to several thousand hours. The authors have investigated the nature of structural change giving rise to this phenomenon. Irradiation was produced by a BSV-1 tube with Mo anode, a current of 10 ma, and a voltage of 45 kv. Irradiation time ranged from 20 to 950 hours. Laue patterns were photographed to trace the structural changes. These were compared with the

Card 1/1

ACCESSION NR: AP4024997

patterns of a crystal not subjected to irradiation. A difference began to appear at 40-50 hours, and irradiated crystals exhibited increased intensity of diffusion maximums and expansion of the Laue spots. These changes became clearer with prolonged irradiation. After 80-100 hours of irradiation, a well-defined asterism appeared on the Laue patterns, a definite indication of separation of the single crystal into separate segments, variously oriented. After about 200 hours, no further reorientation occurred, but the actual time differed for different crystals. It thus appears that the intensity of x-ray reflection weakens because of "fragmentation" in the crystal through irradiation. Decrease in intensity of reflection is due also to radiation defects in the fragments themselves. It is concluded that point defects arise because of the irradiation, and, when the concentration of defects is considerable and their mobility high, they form stable complexes. Changes in the form and size of these complexes at different stages of irradiation explain the changes in intensity of diffusion maximums, the loss of transparency, and the lamination of strongly irradiated crystals. Fragmentation may be related to the "collapse" of these accumulations (complexes) at some definite stage of irradiation and to the formation of dislocation loops, which are free to generate dislocations under proper conditions. Orig. art. has: 2 figures.

Association: Moskovskiy gosudarstvennyy universitet im. M. V. Lomonosova (Moscow State University.)

Card 2/81

ACCESSION NR: APL039393

S/0070/64/009/003/0342/0346

AUTHORS: Telegina, I. V.; Kolontsova, Ye. V.

TITLE: Kinetics of oriented defects in irradiated crystals of LiF

SOURCE: Kristallografiya, v. 9, no. 3, 1964, 342-346

TOPIC TAGS: oriented defect, anomalous scattering, lithium fluoride, neutron bombardment, two dimensional defect, diffraction, radiation defect, radiation effect, defect formation

ABSTRACT: The authors have employed anomalous scattering to study the defect structure of single crystals of LiF after neutron bombardment at an integral flux ranging from  $5 \cdot 10^{16}$  to  $10^{19}$  neutrons per  $\text{cm}^2$  (for fast neutrons). At values of  $3 \cdot 10^{17}$  to  $7.5 \cdot 10^{18}$  neutrons per  $\text{cm}^2$ , two-dimensional defects appear, oriented in the {100} and {111} planes of the initial crystal. These defects are described as two-dimensional zones having scattering capacities differing from average values. The dimensions of these two-dimensional zones change in similar fashion with an increase in integral flux and with an increase in temperature during annealing

Card 1/2

ACCESSION NR: AP4039393

of the crystal (for any particular radiation dose). On the basis of changes in the effects of two-dimensional diffraction during irradiation and during annealing, and also because of similarities in two-dimensional diffraction in deformed and irradiated crystals, it is suggested that the most probable origin of the two-dimensional zones is related to segregations of vacancies in the {100} and {111} planes. In addition to the two-dimensional zones in LiF crystals irradiated by a flux greater than  $3 \cdot 10^{17}$  neutrons per  $\text{cm}^2$ , one may observe defects oriented relative to the initial crystal that create effects of one-dimensional diffraction along {100}. Orig. art. has: 1 figure.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet im. M. V. Lomonosova (Moscow State University)

SUBMITTED: 16Sep63

ATD PRESS: 3081

ENCL: 00

SUB CODE: SS, IC

NO REF SOV: 005

OTHER: 006

Card 2/2



L 2436-66 EWT(1)/EWT(m)/EPF(c)/EPF(n)-2/T/EWP(t)/EWP(b)/EWA(c)  
IJP(c) JD/JW /JG/GG/GS

ACCESSION NR: AT5023806

UR/0000/62/000/000/0264/0267

AUTHOR: Telegina, I. V.; Kolontsova, Ye. V.; Zubenko, V. V.

TITLE: Radiation damage in lithium fluoride crystals

SOURCE: Soveshchaniye po probleme Deystviye yadernykh izlucheni na materialy.  
Moscow, 1960. Deystviye yadernykh izlucheni na materialy (The effect of nuclear  
radiation on materials); doklady soveshchaniya, Moscow, Izd-vo AN SSSR, 1962,  
264-267

TOPIC TAGS: lithium fluoride, irradiation damage, neutron irradiation

ABSTRACT: Neutron-irradiated lithium fluoride single crystals were investigated by means of anomalous x-ray scattering with monochromatized Mo radiation. In crystals bombarded with a total flux of  $(3-6) \cdot 10^{18}$  n/cm<sup>2</sup>, two-dimensional disturbances oriented along the [100] and [111] planes were observed. Annealing of the crystals eliminates these disturbances almost completely, but even after prolonged annealing at 650C, the disturbances oriented along planes of type (100) are preserved. At annealing temperatures above the melting point of lithium, the separation of metallic lithium having a body-centered Bravais lattice is observed in the irradiated crystals. Almost no two-dimensional diffraction effects are observed on x-ray powder patterns of single crystals irradiated with a flux up

Card 1/2



L 2436-66

ACCESSION NR: AT5023806

to  $1 \times 10^{19}$  n/cm<sup>2</sup>. It is concluded that an intense radiation annealing takes place during the irradiation. Orig. art. has: 3 figures.

ASSOCIATION: none

SUBMITTED: 18Aug62

ENCL: 00

SUB CODE: NP, '89

NO REF SOV: 002

OTHER: 015

Card 2/2 *nd*

L 8581-66

ACC NR: AT5023805 EWT(m)/T/EWP(t)/EWP(b) EMA(h)/EMA(c) IJP(c) JD/JG

SOURCE CODE: UR/0000/62/000/000/0257/0263

AUTHOR: Kolontsova, Ye. V.

ORG: none

TITLE: Radiation and deformation defects in crystals

SOURCE: Soveshchaniye po probleme Deystviye yadernykh izlucheni na materialy. Moscow, 1960. Deystviye yadernykh izlucheni na materialy (The effect of nuclear radiation on materials); doklady soveshchaniya. Moscow, Izd-vo AN SSSR, 1962, 257-263

TOPIC TAGS: irradiation, roentgen irradiation, neutron irradiation, plastic deformation, crystal defect, radiation crystal defect, deformation crystal defect

ABSTRACT: The effect of x-ray or neutron irradiation and of plastic deformation on the structure of ionic crystals (LiF, NaCl, KCl, AgCl) and of metal crystals (Al, Zn, Cd, Sn) has been investigated. It was found that both irradiation and deformation generate additional interference patterns which appear on the background of diffusion peaks. As the intensity of irradiation or deformation increases, the number of these patterns increases to such an extent that they form a single arched dash. Under the effect of irradiation or deformation, single crystals break into fragments and their orientation becomes disrupted. The extent of crystal fragmentation depends on the crystal structure and the intensity of irradiation or

Card 1/2

L 8581-66

ACC NR: AT5023805

3

amount of deformation. The changes in LiF crystal structure induced by neutron irradiation and plastic deformation are identical, as proved by the increase in the appearance of diffusion peaks observed during neutron irradiation of diamond, silicon, molybdenum, and quartz single crystals and in the deformation of aluminum single crystals. Since the changes in crystal structure produced by irradiation and deformation are similar, the residual defects in crystals produced by deformation and irradiation are also similar. Orig. art. has: 5 figures. [ND]

SUB CODE: MM,SS/ SUBM DATE: 18Aug62/ ORIG REF: 011/ OTH REF: 005

Card 2/2 p10

KOLONTSOVA, Ye.V.; TELEGINA, I.V.

Structural changes in neutron irradiated  $\alpha$ -quartz. Fiz. tver. tela  
7 no.9:2730-2734 S '65. (MIRA 18:10)

1. Moskovskiy gosudarstvennyy universitet imeni M.V.Lomonosova.

ACC IR: AP6037007

(A, N)

SOURCE CODE: UR/0181/66/008/011/3412/3414

AUTHOR: Kolontsova, Ye. V.; Telegina, I. V.

ORG: Moscow State University im. M. V. Lomonosov (Moskovskiy gosudarstvennyy universitet)

TITLE: Structural variations in quartz during the  $\alpha \rightarrow \beta$  transition and following neutron irradiation

SOURCE: Fizika tverdogo tela, v. 8, no. 11, 1966, 3412-3414

TOPIC TAGS: quartz, phase transition, neutron irradiation, neutron scattering, temperature dependence, atomic property

ABSTRACT: This is a continuation of earlier work (FTT v. 7, 27, 1965) dealing with phase transitions induced in quartz by neutron irradiation. The authors compare the changes as recorded by the diffuse-scattering method or by the Lane method when a quartz is heated, with the structural transformations observed in a quartz following neutron irradiation. The results show that with increasing temperature the temperature  $\alpha \rightarrow \beta$  transition is quite close to the structural changes which are observed in a quartz bombarded with neutrons at fluxes from  $10^{19}$  to  $(5 - 6) \times 10^{19}$  neut/cm<sup>2</sup>. Evidence in favor of this statement is afforded by the gradual nature of the diffuse scattering and by comparisons of the behavior of the Lane maxima and the diffuse

Card 1/2

ACC NR: AP6037007

maxima. The authors interpret the closeness of the structural transitions as a natural result of the fact that the transitions are results of displacements, which should exert the same effect on the atomic fields irrespective of the causes of the displacements. Orig. art. has: 2 figures.

SUB CODE: 20/ SUBM DATE: 25Mar66/ ORIG REF: 005/ OTH REF: 002

Card - 2/2

KOLONTYRSKAYA, Ye.

~~\_\_\_\_\_~~  
A collective's will. Leg.prom. 7 no.11:32-33 H '47.

(MIRA 6:11)  
(Textile industry)



DOKUCHAYEV, V.M., dots.; KOLONUTOV, G., nauchn. red.

[Chemistry in weed control] khimiia v bor'be s sorniakami.  
Stavropol', Stavropol'skoe knizhnoe izd-vo, 1964. 33 p.  
(MIRA 18:8)

1. Stavropol'skiy sel'skokhozyaystvennyy institut (for  
Dokuchayev).

KOLORZ, A.; LOMBERG, K.;

Interface reaction between liquid cast iron and molding material  
with special regard to carbonaceous additions to the molding sand.  
Slevarenstvi 11 no.8/9:374-382 Ag '63.

1. Institut für Giessereitechnik, Düsseldorf.

KOLOS, A.V. (Moskva)

Refining of the classical theory of bending of circular plates.  
Prikl. mat. i mekh. 28 no.3:582-589 My-Je'64 (MIRA 17:7)



KOLOS, A.V. (Lugansk)

Methods for refining the classical theory of flexure and distention  
of plates. Prikl. mat. i mekh. 29 no.4:771-781 J1-Ag '65.

(MIRA 18:9)

KOLOS, A. YE.

KOLOS, A. YE. Abdominal hernias in horses.

So: Veterinariya; 23; 5-6; May/June 1946; Uncl.  
TABCON

KOLOS, A. YE.

KOLOS, A. YE. Post-operative treatment of hemiplegia of pharynx in horses.

So: Veterinariya; 23; (12); December 1946; Uncl.

TABCON



KOLOS, A. V., Sr. Sci. Co-Worker

All-Union Inst. of Exptl. Veterinary Med.

"Intravenous injection of medicinal agents into a horse."

SO: Veterinariia 24(1), 1947, p. 34.

Kolos, A. Ye.

USSR/Diseases of Farm Animals. The Pathology of  
Multiplication

R-3

Abs Jour: Ref Zhur - Biol., No 1, 1959, 2837

Author : Kolos, A. Ye.

Inst : Bashkirian Institute of Agriculture

Title : Data on the Etiology, Pathogenesis, and Treat-  
ment of Postnatal Diseases in Mares

Orig Pub: Tr. Baskirsk. s.-kh. in-ta, 1957, 8, No 2, 415-  
426

Abstract: No abstract

Card 1/1

Distr: 4E3b/4E2c(j)

✓ The reactivity of some ionized hydrocarbons. B. Kolos,  
(Inst. Badań Jądrowych, Warsaw). *Polish Acad. Sci.  
Inst. Nuclear Research, Rept. No. 119/V, 10 pp. (1955) (in  
Russian)*.—Charge distribution, free valencies, and bond  
orders for neutral and ionized mols. of butadiene, hexa-  
triene, anthracene, 1,2-benzopyrene, and 1,2,3,4-dibenzo-  
pyrene are given and discussed. A. Szafranski

SR

3  
1-92 (NA)  
2

KOLOS, E.

✓ Assay of frangula-bark extract. S. Simon and E. Kolos  
(Magyar Pharm., Budapest). *Acta Pharm. Hung.* 5, 195-203 (1954) in German. — The acetone-pptd.,  
concd. MeOH ext. contained more total cincholin than ace-  
tone-pptd. aq. aq. EtOH, or MeOH frangula-bark ext. In 2  
instances official Pharm. exts. had little biol. activity.  
There was no parallelism between the results of biol. and  
chem. assays.  
S. Ellis

KOLCS, Ede, a kémiai tudományok kandidátusa

The state and analytic chemical problems of manufacturing fine chemicals in Hungary. Kem tud kozl 20 no.3:323-326 '63.

1. Reanal Finomvegyszergyár, Budapest.

KOLOS, F.;WIEDNER, L.

KOLOS, F.;WIEDNER, L.

Completed greasing experiments with domestic sulfonated oils on chrometanned leathers. p. 81

Vol. 5, No. 4, August, 1955 Budapest, Hungary BOR-E5 CIPOTECHNIKA

SO: Monthly List of East European Accessions, (EEAL), IC, Vol. 5  
No. 3, March, 1956

KOLOS, F.

New Methods in the field of liming, p. 1. BORO ES CIPOTECHNIKA. (Boripari Tudományos Egyesület mint a Magyar Tudományos Egyesületek Szövetsége Tagegyesülete) Budapest. Vol 6, no. 1, Feb. 1956

SOURCE: East European Accessions List (EEAL), Library of Congress Vol 5, no. 6 June 1956

KOLOS, Imre, foallatorvos (Sukosd); RETI, Pal (Budapest V., Rizenberg hp.u.17)

Motorists' letters. Auto motor 13 no.21:8 1 N '60.



KOLOS, I.

In an Azov machine repair station. Sov.profsoiuzy 6 no. 9:54-56  
Ag '58. (MIRA 11:8)

(Repair and supply stations)

KOLOS, I.

Let's reorganize the remuneration of rural machinery operators.

Sov. profsoiuzy 6 no.12:54-56 S '58.

(MIRA 11:9)

(Collective farms--Accounting)

KOLOS, I.

Integrated approach. Sov.profsoiuzy 6 no.14:23-25 0 '58.  
(MIRA 11:12)

1. Sovkhoz imeni Stalina Skvirskogo rayona, Kiyevskoy oblasti.  
(Kiev Province--State farms)

KOLOS, I.

Kuban, land of oil. Sov.profsoiuzy 7 no.2:34-37 Ja '59.

(MIRA 12:3)

(Kuban--Oil well drilling)

KOLOS, I.

New page in the life of a collective-farm village. Sov.  
profsoiuzy 7 no.4:45-48 Fe '59. (MIRA 12:5)  
(Ivanovskoye--Agricultural laborers)

KOLOS, I.A.; ORLOV, V., red.

[People of the great flight] Liudi bol'shogo poleta. Moskva, Politizdat, 1965. 117 p. (MIRA 18:7)

PAL, Mihaly (Budapest); KULOS, Janos (Budapest)

forum of innovators. Ujit lap 16 no.14:30-31 25 JI '64.



KOLODS, L.

✓ 768. Pneumatic valve for inflatable members.  
 "LICENCIA" TALLERES Y FABRICAS VALLA-  
 BAT, inv. L. Kolod. U.S.P. 2710623; appl. 17.1.52; NT  
 Hungary 4.7.51; sec. 14.6.55. A valve for bladders  
 of balls, inner tubes of tyres, and the like, comprises  
 an integral body of resilient material having a  
 longitudinal central portion and a pair of flanges  
 formed at opposite ends of the central portion and  
 defining a recess between them to receive a rein-  
 forcing member around the central portion, and a  
 passage extending through the flanges and central  
 portion, this passage being defined by a pair of flat  
 slits extending in intersecting planes, narrowing  
 towards each other and meeting at the meeting point  
 of the adjacent apices of the triangular conforma-  
 tions defining the slits. 66Q31:313.1

(1)

KOLOS, L.A., assistant

Permeability to air of a double-layer sleeve in a specific case.  
Nauch.trudy MTILP no.23:150-154 '61. (MIRA 15:9)

1. Kafedra teplotekhniki Moskovskogo tekhnologicheskogo instituta  
legkoy promyshlennosti.

(Coats)

KOLOS, L.A., assistant

Wear of the heel in multiple repeated loading. Nauch. trudy MTILP  
no.29:150-151 '64. (MIRA 18:4)

1. Kafedra teplotekhniki Moskovskogo tekhnologicheskogo instituta  
legkoy promyshlennosti.

KOLOS, L.A., starshiy prepodavatel'

Lead action on the heels. Nauch. trudy MTILP no.30:136-139 '64.

Impact load on heels. Ibid.:140-142

(MIRA 18:6)

1. Kafedra teplo tekhniki Moskovskogo tekhnologicheskogo instituta legkoy promyshlennosti.

KOLOS, L.A., assistant

Heel wear. Nauch. trudy MTILP no.27:173-174 '63.

(MIRA 17:11)

1. Kafedra teplostechniki Moskovskogo tekhnologicheskogo instituta  
legkoy promyshlennosti.

KOLOS, N.D., inzh.; VORONCHIKHIN, G.P., inzh.

Simple method for preparing metal surfaces fo. painting.

Mashinostroenie no.1:76-78 Ja-F '65.

(MIRA 18:4)

Kolos, P.

197. Stuffing experiments with domestic sulphonated oils in chrome-tanned leathers. P. Kolos, L. Wiedner, *Ist. Isipiterchika*, Vol. 5, 1955, No. 4, pp. 81-86, 7 figs., 4 tabs.

Stuffing experiments had been conducted on leathers where the circumstances of manufacture and their applications differed greatly namely chrome-side leather made from native hides and glove leather. For chrome-side leather a standard liquoring method with sulphonated sunflower-seed oil was applied. It was

found that from the available oils rapeseed oil is more adequate for liquoring purposes than sunflower-seed oil. The best results were obtained with rapeseed oil having a sulphonation degree of  $19 \pm 1$ . Thus it was possible to reduce the quantity of grease used by 20% and furthermore the obtained leather possessed better properties (closed grain, smaller moulder-side, elastic and soft touch). In spite of some statements in literature which claim that sulphonated train oils are best it was found that sulphonated neats' foot oil was superior for stuffing glove leather from pig skin bellies. Fatty spears do not necessarily occur in stuffing with neats' foot oil; this phenomenon is primarily due to incorrect sulphonation or to the oxidized fatty acids enriched by excessive sulphonation. It could be established that sulphonated neats' foot oil with a sulphonation degree of  $19 \pm 1$  produced the best results if the sediments consisting principally of oxidated fatty acids do not exceed 5% by volume.

Maths

2

KOLOS, P.

Tracks on sectionalized prestressed-concrete ties with concrete slits.

p. 213 (Przegląd Kolejowy Drogowy. Vol. 8, no. 9, Sept. 1956. Warszawa, Poland)

Monthly Index of East European Accessions (EEAI) LC. Vol. 7, no. 2,  
February 1958



Kolos, P. I.

USSR / Soil Science. Cultivation. Melioration. Erosion. J-5

Abstr Jour: Ref Zhur-Biol., No 8, 1958, 34433.

Author : Panenko, I. D., ~~Kolos, P. I.~~

Inst : Moldavian Experiment Station for Irrigation of  
Vegetables and Potatoes.

Title : Importance of Power Irrigation for the Increase  
of Yield of Agricultural Cultivations.

Orig Pub: Tr. Mold. ovosheho-kartof. orosit. opyt. st.  
Kishinev, Gosizdat Moldavii, 1956, 77-92.

Abstract: Power irrigation permits a better utilization of  
moisture, and is more effective in agricultural  
cultivations, increasing the yield as follows:  
apple trees - by 16-50%, viticulture - by 23-35%,  
tomatoes by 37%, potatoes by 41%, and that of  
winter wheat on black fallow by 36-90%. The high-  
est yields obtained, were those in combined power

Card 1/2

Kolosa T.

Watering of vineyards in southern part of the Dnieper Valley. P. I. Kolosa, Solonchik, Finogradovo, P. I. de la Madaia II, No. 4, 19-20/10/57. Watering of vineyards in the region 3 times during the summer season at 700 cm.m. water, for treatment of the vines by 10-17 centners/ha. without affecting the grape quantities (sugar 13.16-22.72%; total acidity 0.48-0.63 g/l. must, also content of the wine 12.7-13.8%). E. Wierbicki

KOLOS, R.

TECHNOLOGY

PERIODICAL: MAGYAR HIRADASTECHNIKA. Vol. 9, no. 2/3, June 1958

Kolos, R. The results of the Exhibition of Telecommunication Engineering and the third Conference of Telecommunication Engineering. p.45.

Monthly List of East European Accessions (EEAI) LC, Vol. 8, No. 2,  
February 1959, Unclass.

KOLOS, Richard, egyetemi tanar; FISCHER, Imre; HARSANYI, Istvan, dr., docens, kozgazdasagi tudomanyok kandidatusa; LENGYEL, Karoly, dr., tudomanyos munkatars; BALASSA, Laszlo, foeloado; KOZMA, Pal, dr., fokonyvelo; KASPER, Egon, dr.; MACSKASZV, Pal; LUKACS, Laszlo, okleveles elektromernok; SANTA, Balint, dr.

Scientific conference on retraining instrument industry personnel.  
Meres automat 13 no.2/3:66 '65.

1. Secretary General, Council on Science and Higher Education, Budapest (for Kolos).
2. Head, Directorate of Instrument Industry of the Ministry of Metallurgy and Machine Industry, Budapest (for Fischer).
3. Instrument Industry Research Institute, Budapest (for Lengyel).
4. National Price Office, Budapest (for Balassa)
5. Instrument and Business Machine Trading Organization, Budapest (for Kozma).
6. Group Head, Ministry of Finance, Budapest (for Kasper).
7. Directorate of Instrument Industry of the Ministry of Metallurgy and Machine Industry, Budapest (for Santa).

KOLOS, Richard, tanszekvezeto docens

National Conference on Instrumentation; presidential opening address.  
Meres automat 8 no.9:257-258 '60.

1. Miniszterhelyettes, es Merestechnikai es Automatizalasi Tudomanyos Egyesulet elnoke, Budapest.

(Hungary--Measuring instruments)

KOLOS, Richard, tanszekvezeto docens

Tasks of the Scientific Association of Measuring Technique  
and Automation. Meres automat 9 no.2:33-36 '61.

1. Miniszterhelyettes, es Merestechnikai es Automatizalasi Tudos-  
manyos Egyesulet elnoke

KOLOS, Richard, miniszterhelyettes, tanszékvezetődocens

The chairman's opening address to the National Conference on Instrumentation. ~~Meres automat~~ 8 no.9:257-258 '60.

1. A MATE elnöke.

KOLOS, Richard, egyetemi tanar

Basic research and production. Elet tud 17 no.47:1483-1484  
25 N '62.

1. Tudomanyos es Felsoktatasi Tanacs fotitkara.



HEROUT, V.; KOLOS, F.; FLIVA, J.

Terpenes. Part 49. Sesquiterpenes of the cadinene type in Javanese citronella oil [abstract; in English]. Sbor.Chekh.khim.rab. 18 no.6: 886 D '53. (MLRA 7:6)

1. Department of Natural Substances, Institute of Organic Chemistry, Czechoslovak Academy of Science, Prague. (Cadinene)

KOLCS, T.

HEROUT, V., KOLCS, T., PLIVA, J.

"Terpenes. Part 49. Sesquiterpenes of the Cadinene Type of Javanese Oil of Citronella," p. 440.

(Chemicke Listy, Vol.47, No.3, Mar. 1953, Praha.)

SO: Monthly List of East European Accessions, Vol.2, No.9, Library of Congress, September 1953, Uncl.

AKIMENKO, I.S.; KOLOS, T.K.; MERKIN, V.G.; SMOTRICH, B.A.; YASENSKAYA, M.T.

Method of water-and-heat treatment of corn. Fern. i spirt.prom.  
31 no.3:36-37 '65. (MIRA 18:5)

1. Lipetskiy spirtozavod.

BALON, I.D., kand.tekhn.nauk; ROMANENKO, N.T., inzh.; YUPKO, L.D., inzh.;  
BOLKUNOV, Ye.P., inzh.; TULUYEVSKAYA, T.A., inzh.; ASTAFUROV, P.I., inzh.;  
VOLOVIK, A.V., inzh. Priniimali uchastiye: BAKAYEV, A.I.; VOKHNIK, A.R.;  
KOLOS, V.D.; KAYSTRO N.P. [deceased]; LITVINENKO, V.I.; MAKARCHENKO, N.M.;  
ONOPRIYENKO, V.P.; PALAGUTA, V.P.; PIKA, V.S.; RAGIN, B.I.; ROMANCHENKO,  
Ye.I.; SAYENKO, S.D.; STOLYAR, V.V.; SKORIK, N.M.; TOROPENKO, P.D.

Characteristics of making ferromanganese in large capacity blast furnaces  
and the effect of slag conditions on basic technical and economic indices.  
Stal' 23 no.12:1069-1073 D '63. (MIRA 17:2)

1. Ukrainskiy nauchno-issledovatel'skiy institut metallov i zavod "Zapo-  
rozhtal".

24 (6)

AUTHORS: Severdenko, V. P., Academician, AS BSSR, SOV/20-126-5-14/69  
Kolos, V. I.

TITLE: On a Field of Slide Lines (Ob odnom pole liniy skol'zheniya)

PERIODICAL: Doklady Akademii nauk SSSR, 1959, Vol 126, Nr 5, pp 964 - 965  
(USSR)

ABSTRACT: The solution of practical problems arising from the theory of plasticity frequently involves fields of slide lines similar to that shown by figure 1. In this case complicated numerical problems permit a solution of the problem; some particularities, however, offer a less complicated and more exact solution. A field is discussed in which families of lines originate from two centers  $O_1$  and  $O_2$ , which intersect each other. The arcs originating from  $O_1$  are assumed to have a common radius of curvature  $R$ , those originating from  $O_2$  have a radius of curvature  $S$ ,  $m$  lines originate from  $O_1$  within the total angle  $\alpha$ , and  $n$  lines from  $O_2$  within the total angle  $\beta$ . These (curved) lines are separated from one another at  $O_1$  by the angle  $\Delta\alpha_1$  and at  $O_2$

Card 1/2

On a Field of Slide Lines

SOV/20-126-5-14/69

by the angle  $\Delta\beta_j$  (Fig 1). The authors then set up the line of the nodes  $R_{m,n}$  and  $R_{\alpha,\beta}$ , which are transformed for an (xy)-coordinate system. Further, a formula is given for the simplified case in which  $\alpha = \beta$ . The authors evaluated the results numerically for a 15-degree net of lines, and the coordinates of the nodes, including the arcs of  $135^\circ$ , are listed in a table. The radii of the nodes are computed by the infinite series  $R_{\alpha,\beta} =$

$= 1 + \frac{\alpha}{1!} + \frac{\alpha\beta}{1!1!} + \frac{\alpha^2\beta}{2!1!} + \frac{\alpha^2\beta^2}{2!2!} + \frac{\alpha^3\beta^2}{3!2!} + \dots$ . There are 1 figure, 1 table, and 2 Soviet references.

SUBMITTED: April 2, 1959

Card 2/2

KOLOS, V.I. (Volgograd)

Insertion of concave stamp into a plastic medium. PMTF no.2:157  
Mr-Ap '65. (MIRA 18:7)

KOLOS, V. I.

Cand Tech Sci - (diss) "Force and performance of deformation in stamping." Minsk, 1961. 11 pp; (Academy of Sciences Belorussian SSR, Technical Physics Inst); 100 copies; price not given; (KL, 5-61 sup, 190)



KOLOS, V.I.; TRET'YAKOV, Ye.M.

Determining the thickness of a solidified layer. Kuz.-shtam.  
proizv. 5 no.2:47-49 P '63. (MIRA 16:2)  
(Thickness measurement)

KOLOS, V.I. (Volgograd)

Integrating equations for the plane deformation of a totally  
plastic body. Izv. AN SSSR. Otd. tekhn. nauk. Mekh. i mashinostr.  
no. 1:171-173 Ja-F '63. (MIRA 16:2)  
(Deformations (Mechanics)) (Plasticity)

S/179/63/000/001/025/031  
E081/E135

AUTHOR: Kolos, V.I. (Volgograd)

TITLE: Integration of the plane deformation equations of an ideally plastic body

PERIODICAL: Akademiya nauk SSSR. Izvestiya. Otdeleniye tekhnicheskikh nauk. Mekhanika i mashinostroyeniye, no.1, 1963, 171-173

TEXT: The paper deals with the integration of the equations  
 $dR - Sd\alpha = 0, dS + Rd\beta = 0, du - v d\beta = 0, dv + u d\alpha = 0 \quad (1.2)$

where  $R, S$  are the radii of curvature of the characteristics;  $u, v$  are the velocities along the characteristics;  $\alpha, \beta$  are angles defined as in Fig.1. These equations are developed into infinite series by successive substitution. The series are evaluated for characteristics formed by circular arcs and the results tabulated for  $15^\circ$  intervals of  $\alpha$  and  $\beta$ . The characteristics arising from the deformation of a plastic cylinder by rigid cylinders are evaluated and illustrated, and the case of a plastic region contained between moving rigid regions is also discussed.

Card 1/2


The integration of the plane ...

S/179/63/000/001/025/031  
E081/E135

There are 3 figures and 1 table.

SUBMITTED: August 1, 1962

Fig. 1



Card 2/2

KOLOS, W.

Theoretical examination of the dibenzo-derivatives of perylene. In English.

p. 3, (ACTA UNIVERSITATIS SZEGEDIENSIS) Vol. 2, no. 1/4, 1956  
Szeged, Hungary

SO: Monthly Index of East European Accessions (EEAI) LC, Vol. 7, No. 3,  
March 1958

KOLDS, Walter.

Distr: 4E3d/4E2c(j)

43. A theoretical study of the dibenzo derivatives of perylene. P. BERENCSE, W. KOLDS, R. PAUTECZ. *A Magyar Tudományok Akadémiájának Kémiai Tudományok Osztályának Közlekedései*. Vol. 9, 1957, No. 3, pp. 293-298, 2 figs.

4  
2 May  
2

The ternary systems of aromatic hydrocarbons having condensed rings consisting of anthracene or phenanthrene parts have been investigated by means of the I. C. A. O.

method and comparative calculations have been made as to the practicability of perturbation computations. The calculations resulted in a qualitatively true representation of the principal characteristics of the spectra but quantitative agreement is still not satisfactory.

gjd

Kolos, W.


3  
IRML

1071  
Kolos W. The Influence of Hindered Rotation on the Scattering of  
Fast Neutrons by Bound Protons.

A. A. Physics, Poltava (Ukraine).  
The cross section of the proton-neutron  
interaction depends on the degree of freedom of the  
proton. The cross section of the proton-neutron  
interaction is calculated for a hindered rotation of the  
proton in a molecule. It is shown that the cross section  
of a proton bound in a molecule is smaller than  
that of a free proton. It is also shown that the  
cross section of a proton bound in a molecule is  
smaller than that of a free proton.

"APPROVED FOR RELEASE: 09/18/2001

CIA-RDP86-00513R000823920014-2



APPROVED FOR RELEASE: 09/18/2001

CIA-RDP86-00513R000823920014-2"



115341

The type of proton considered is a molecule which prefers to rotate in a direction with respect to the  $z$ -axis. The sections are calculated for a molecule in which rotational oscillations in the  $z$ -direction are excited or de-excited. In the scattering of slow neutrons by methyl alcohol is discussed.

KELSON

3

POL

539,185

4380. The influence of hindered rotation on the scattering of slow neutrons by liquid paraffins. W. Kozos. *Bull. Acad. Polon. Sci. Cl. 5, 2, No. 9, 433-4* (1954).

Calculations are presented of the molecular cross-section of methyl alcohol, including the effect of inelastic scattering of the neutrons with excitation of rotational levels. The results do not agree well with experiment, and it is suggested that this is largely because the effect of molecular vibration was neglected.

J. M. CASSELY

EmL 2/11

Kalcs Włodzimierz

6929. The influence of hindered rotation on the

viscosity of polymer solutions

1964

KOLOS, W.

Tribute to Einstein. p. 269. Vol. 1, no. 3, 1955      Warszawa

SERIA B: PRZYROD A NEOZYWIONA

SOURCE:      East European Accession List (EEAL) Library of Congress  
Vol. 5, no. 8, August 1956

KOLOS, W.

B-4

Category: Poland / Physical Chemistry - Molecule. Chemical bond.

Abs Jour: Referat Zhur - Khimiya, No 9, 1957, 29627.

Author : Kolos Wlodzimierz

Inst : not given

Title : The influence of Hindered Rotation on the Scattering of Slow Neutrons by Bound Protons. II.

Orig Pub: Acta phys. polon., 1955, 14, No 3, 173-182

Abstract: An investigation of the effect of hindered rotation in the molecule on cross section of neutron scattering by proton bound within the molecule. Previously (Communication I, RZhKhim, 1955, 54457) there was derived the formula for the cross section  $\sigma_{mf,mpi} = b_0^2 \sigma_{mf,mpi}$  wherein  $\sigma_{mf,mpi}$  is cross section of scattering of neutrons at rigid molecule, taking into account the rotation. In the present paper is proposed a procedure of summation, by states of different  $mf$ , that is suitable for energy of neutrons, comparable with distances between rotation levels in the molecule. Use of the static approxima-

Card : 1/3

-20-

Polish Acad Sci, Warsaw.

APPROVED FOR RELEASE: 09/18/2001

Category: Poland / Physical Chemistry - Molecule. Chemical bond.

Abs Jour: Referat Zhur-Khimiya, No 9, 1957, 29627

tion of Placzek (Placzek G., Phys. Rev., 1952, 86, 377) permits to disregard change in energy of neutron, associated with transitions within the molecule, which simplifies summation by states in the cross section formula. Calculations are performed at Born approximation. As psi-functions are chosen the eigenfunctions of symmetrical rotator:  $U_{m,m'}(\theta, \psi, \varphi) = \sqrt{\frac{1}{2} \pi} \times \frac{(2l+1)}{2} \sqrt{\frac{1}{2} \pi} e^{-im\theta} e^{-im'\psi} e^{-im\varphi} \exp \left[ \frac{1}{2} (1 + i) \frac{m}{l} \right] Q_m = t^{l/2} (1-t)^{l/2} \times F(l, m, t)$ , where  $F$  -- hypergeometric series. Interaction of neutrons with the molecule is defined by the known Fermi potential  $V = \sum m_i \delta(r - r_i)$ . Finally the cross section is written in the form:

$$\sigma_m = \frac{4 \pi b_0^2 (1 + 2 \nu 3^{\nu})^{1/2}}{(2l+1)! (2l+3)!} \times \sum_{m'} \sum_{n=0}^{\infty} (-1)^{m'} D_n +$$

wherein  $1, 2, 3^{\nu}$  -- principal values of reduced tensor of molecule masses;

$$I_m = \int_0^1 t^{\nu} (1-t)^{\nu} F^2(m, t) dt, \text{ and } D_n \text{ -- number series.}$$

Card : 2/3

-21-

STATISTICAL  
1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218  
2219  
2220  
2221  
2222  
2223  
2224  
2225

been calcd from the theory presented in  
48. 8672e; 49, 9370A). The effect of rotation, modeled  
on the variation on the cross-section is taken into  
to motion, or the effect of rotation on the  
between the calcd. and exptl. values should be a measure of its  
contribution. However, this effect would be shown  
strongly by a motion like Model, also  
has been partially ruled, previously (cf. C.A. 49, 9370A).

POLAND/Physical Chemistry - Molecule. Chemical Bond.

B-4

Abs Jour : Referat Zhur - Khimiya, No 6, 25 March 1957, 18203

Author : Kolos

Inst :

Title : Effective Cross-Section of the Molecule of Methyl Mercaptan for Dispersing Slow Neutrons.

Orig Pub : Bul. Polskoy AN. 1956, Otd. 3, 4, No 5, 263-265

Abstract : Work on computation of an effective cross-section of molecules for dispersing slow neutrons is continued (RZhKhim, 1955, 54457). For computation of proton effective cross-section the association of molecules in liquid state is taken into consideration. Intramolecular bonds in associated complexes can hinder a free rotation of molecules which results in increasing effective cross-section of protons in comparison with the case of a free rotation of molecules. This was not taken into consideration in previous work where the difference between a computed and an

Card 1/4

- 37 -

where

Card 2/4

- 38 -

APPROVED FOR RELEASE: 09/18/2001

CIA-RDP86-00513R000823920014-2

state, respectively. In this formula there is a typographical error in the previous work. For computation the height of the potential barrier was taken equal to

Card 3/4

- 39 -

Abs Jour : Referat Zhur - Khimiya, No 6, 25 March 1957, 18203

705 cal/mol. As the energy of the bond O-H...O is considerably greater than the energy of S-H...S, the association is small in case of CH<sub>3</sub>SH in comparison with the case of CH<sub>3</sub>OH. Therefore, the agreement of the experimental and computed effective cross-section for CH<sub>3</sub>SH will mean that the influence of the association upon effective cross-section does not exist and that the pre-supposed appraisal of the influence of vibrations of protons upon the effective cross-section is correct. Curves of the dependence of effective cross-section of the molecule CH<sub>3</sub>SH are given. For computations the height of potential barrier is taken equal to 705 cal/mol without and with taking into consideration the influence of vibrations of protons as a function of the initial energy of the neutron.



~~KOLOS, W.~~

"A certain connection between science and politics."

p. 15 (Kosmos, Serbia B: Przyroda Nieożywiona) Vol. 3, no. 1, 1957  
Warsaw, Poland

SO: Monthly Index of East European Accessions (EEAI) LC. Vol. 7, no. 4,  
April 1958

KOLOS, W.  
KOLOS, W.

On the one-center interaction integrals in the molecular orbital method.

p. 299 (Acta Physica Polonica) Vol.15, no. 3, 1957, Warszawa., Poland

SO: MONTHLY INDEX OF EAST EUROPEAN ACCESSIONS (EEAI) LC, VOL. 7, NO. 1, JAN. 1958

KOLOS, WLODZIMIERZ  
APPROVED FOR RELEASE: 09/18/2001

CIA-RDP86-00513R000823920014-2

POLAND/Atomic and Molecular Physics - Physics of the Molecule

Abs Jour : Ref Zhur - Fizika, No 6, 1958, No 12871

Author : Kolos Wlodzimierz  
Inst : Polish Academy of Sciences, Warsaw  
Title : Electron Correlation in Molecules.I. The Ethylene Molecule

Orig Pub : Acta phys. polon, 1957, 16, No 4, 257-266

Abstract : As is known, calculation of the electron correlation in the molecular-orbit method leads in general to great difficulties. In connection with this, the author proposes a method for partial accounting for the electron correlation, which takes into account only the electrostatic correlation of the electrons with oppositely-directed spins, belonging to one pair, and neglects the electrostatic correlation of the electrons with like spins and of electrons with opposite spins but belonging to different pairs. The method can be used successfully for the calculation of the excitation energy of conjugated hydrocarbonates and it is illustrated with ethylene as an example. Using  $\pi$ -electron wave functions in the interaction

Card : 1/2

Card : 2/2

KOLOS W.

POLAND/Atomic and Molecular Physics - Physics of the Molecule.

D

Abs Jour : Ref Zhur Fizika, No 8, 1959, 17569

Author : Kolos, Wlodzimierz

Inst : Institute of Physics, Polish Academy of Sciences, Warsaw  
Poland

Title : On the Electron Density Distribution in Molecules.

Orig Pub : Acta phys. polon., 1958, 17, No 2-3, 201-202

Abstract : General ideas are expressed concerning the character of the distribution of the electron density in simple molecules. The wave function of the system is represented in the form of a product of Slater single-electron atomic orbits and a correction factor, which takes into account the neutral perturbation; its form i.e., indeed determined by the electron density distribution in the molecule. -- K.P. Gurov

Card 1/1

- 45 -

19

APPROVED FOR RELEASE: 09/18/2001

Distr: 4E2c(j)/4E3d

✓ A method of interpretation of absorption spectra of complex ions. Wlodzimierz Kolos (Polish Acad. Sci., Warsaw). *Roczniki Chem.* 32, 329-38 (1958) (English summary).—The H-like atom with one electron is considered to account for the electronic configurations of various atoms. The latter may be found by solution of Schrödinger's equation, giving a set of corresponding at. orbitals, and by filling the orbitals successively with electron pairs. Similarly, the solution for one-electron atom in an outer sym. field was used to explain the features and origin of absorption spectra of complex atoms by taking the results of Ilse and Hartmann (*C.A.* 46, 7874e) and of Ballhausen (*C.A.* 49, 9383b). The 5-fold degenerated d state splits up by action of outer field, depending on the symmetry of perturbation. The split levels were filled with electrons, allowing for the exclusion principle, with the assumption that the perturbation did not change the multiplicity of the ground state. The excitation energies of some complex ions were calcd. by the use of empirical parameters (one for  $O_h$  as in  $Co^{++}$  complexes, and another for  $D_4$  symmetry as in  $(Ni.aq.)^{++}$ ) and found to agree reasonably with exptl. data. In the case of  $Cu^{++}$  complexes the values are identical with those resulting from the crystal field theory. A. Kreglewski

JB

1/

5  
2-MAY  
2

yes

POLAND/Solid State Physics - Solid State Theory

E-2

Abs Jour : Ref Zhur - Fizika, No 1, 1959, No 790

Author : Kolos Wlodzimierz  
Inst : Electrochemical Division Institute of Chemical Physics,  
Polish Academy of Sciences, Warsaw, Poland  
Title : Stability of Certain Complexes Having a Coordination Number  
of 4.

Orig Pub : Roczn. chem., 1958, 32, No 2, 393-395

Abstract : The theory of crystalline fields (in the strong-field approximation) is applied to the calculation of the relative stability of the quadratic and tetrahedral complexes Cu (II), Ni (II), and Co (II). It follows from the calculations that from among the tetrahedral, ones the most stable are the complexes Co (II), while the complexes Ni (II) are the most stable among the quadratic ones. These results are in agreement with the experimental data.

Author's resume

Card : 1/1

82709

P/046/60/005/001-2/003/008  
A222/A026

5.3100

AUTHOR: Kołos, Włodzimierz

TITLE: On the Reactivity of Certain Ionized Hydrocarbons

PERIODICAL: Nukleonika, 1960, No. 1-2, pp. 73-80

TEXT: Charge distributions, bond orders and free valencies have been calculated for butadiene, hexatriene, naphthalene, anthracene, 1,2-benzopyrene and 1,2-3,4-dibenzopyrene. The corresponding molecular diagrams are presented in Fig. 1-3, where a) refers to the ionized and b) to the neutral molecules. The author hopes that the results may be helpful in understanding some elementary processes occurring in radiosyntheses. The calculus refers to the theory of molecular orbitals in the Hückl approximation. Atomic orbital overlap integrals have been ignored. Molecular orbitals for electrons thus were assumed as

$$\Psi_i = \sum_r c_{ir} \Psi_r$$

(1)

Card 1/3

82709

P/046/60/005/001-2/003/008  
A222/A026

# On the Reactivity of Certain Ionized Hydrocarbons

where  $\psi_r$  is the  $2p_z$  orbital of the  $r$ -th carbon atom, and the indicator  $r$  skips all carbon atoms belonging to the system of conjugate bonds. The coefficients  $c_{ir}$  respond the condition of normalization

$$\sum_r |c_{ir}|^2 = 1 \quad (2)$$

The charge at the atom  $r$  originating from the  $\pi$  electrons is defined by the formula

$$q_r = \sum_i m_i |c_{ir}|^2 \quad (3)$$

where  $m_i$  is the number of electrons (in the molecule), whose condition is described by the  $\psi_i$  function. Further calculus applicable for alternant hydrocarbons (all compounds dealt with in the article) is cited from references 1 through 5. It has been proved that in ionized polyenes the largest electric charge must be located at the ends of the chains. Proof has been found for a smoothing effect of the ionization on bond orders and an increase in free valencies of carbon atoms at the ends of

Card 2/3

82709

P/046/60/005/001-2/003/008  
A222/A026

On the Reactivity of Certain Ionized Hydrocarbons

the molecules. The largest electric charge in ionized naphthalene and anthracene molecules must be located in the  $\alpha$  and mezo positions respectively, and the smallest charge in the  $\beta$  positions. Apparently, the points most susceptible to attack in reactions with electrophilic and nucleophilic reagents are determined by these charge distributions. There are 3 figures and 7 non-Soviet references.

ASSOCIATION: Instytut Badań Jądrowych PAN, Pracownia Chemii Radiacyjnej (Institute of Nuclear Research, PAN, Laboratory of Radiation Chemistry), Warsaw, Instytut Chemii Fizycznej PAN, Zakład Elektrochemii (Institute of Physical Chemistry, PAN, Department of Electrochemistry), Warsaw

SUBMITTED: December 1959

Card 3/3

27316  
P/046/60/005/011/005/018  
D249/D303

21.6000

AUTHOR: Kołos, Włodzimierz

TITLE: Ionization of molecular gases by  $\beta$ -particles

PERIODICAL: Nukleonika, v. 5, no. 11, 1960, 719 - 726

TEXT: The average energy,  $W$ , required to produce an ion pair exceeds the ionization potential,  $I$ , to an extent depending on the ratio of the probabilities of excitation and ionization, since a part of  $W$  merely raises the molecule into a higher energy level. A brief summary of Western work in this field is given. Contrary to early experimental indications, the theory of excitation and ionization of atoms by fast particles predicts a correlation between  $W$  and  $I$ , and shows a dependence of  $W$  on the generalized oscillator strengths for the excitation of a given atom. The relationship between  $W$  and  $I$  has recently been confirmed by experiment, being nearly linear for the inert gases. For the same value of  $I$ , the values of  $W$  are appreciably lower for the monatomic than for the di-

Card 1/5



27316

P/046/60/005/011/005/018  
D249/D303

Ionization of molecular ....

atomic gases. In the present work, hydrogen is used to indicate, on theoretical grounds, that the average energy required to produce an ion pair by  $\beta$ -particles should be greater for diatomic than for monatomic gases. The total cross-section for inelastic collisions is divided into (1)  $\sigma_e$ , corresponding to inelastic collisions leading to the excitation of discrete levels with an average excitation energy  $E_e$ , and (2)  $\sigma_{11}$  and  $\sigma_{12}$ , corresponding to inelastic collisions which lead to ionization and produce secondary electrons with kinetic energies  $E_{11}$  and  $E_{12}$ , where  $E_{11} < I < E_{12}$ . Assuming  $W$  to be independent of the energy of impinging electrons: X

$$W = \frac{\sigma_e E_e + \sigma_{11} (I + E_{11}) + \sigma_{12} I}{\sigma_{11} + \sigma_{12}} \quad (1)$$

Using this equation and the cross-sections calculated by H. Bethe (Ref. 6: Handbuch der Physik. Bd. 24 T.I. Berlin 1933, Springer Verlag) for the various excitations in H-atoms by the first born

Card 2/5

27316  
P/046/60/005/011/005/018  
D249/D303

Ionization of molecular ...

approximation, the following table is constructed:

Table 1. Energy of primary electrons, in kV	100	10	1	0.2
W, in eV per ion-pair	36	34.5	33	27

For hydrogen, up to 50 % of ionization is due to secondary electrons with low energies, for which the Born approximation is not valid. From the above table, W appears to be energy dependent, although this may not be true when all the energies of primary and secondary electrons and the energy distribution are taken into account. Experimental values of W for molecules disagree with those calculated for free atoms by Eq. (1), probably due to the difficulty of calculating precisely the cross-sections for various inelastic collisions. Collision of an electron, with an H<sub>2</sub> molecule is considered, which raises the molecule from the ground to the excited state and the change of momentum is assumed to be small. Using wave mechanics, it is shown semi-qualitatively that the formation

Card 3/5

27316

P/046/60/005/011/005/018  
D249/D303

Ionization of molecular ...

of a diatomic molecule increases the cross-section for the excitation of discrete energy levels ( $\sigma_e$ ). It is pointed out that little is known about molecular wave functions for excited states and that the transition probabilities are very sensitive to the form of wave functions employed. Only a very accurate calculation could, therefore, give conclusive results. Experimental values of the ionization cross-sections, obtained by Western authors for H-atoms, agree with the theoretical ones for energies  $> 250$  ev, but for lower energies the theoretical values are significantly higher. The experimental values for the  $H_2$  molecule are available for comparison with those of 2 H-atoms and tend to support the author's conclusions. There are 1 figure, 1 table and 10 non-Soviet-bloc references. The 4 most recent references to the English-language publications read as follows: W.P. Jesse, J. Sadauskis, Phys. Rev. 90, 1120, 1953; T.E. Bartner, G.S. Hurst, Phys. Rev., 93, 1236, 1954; W.C. Barber, Phys. Rev., 97, 1071, 1955; J. Weiss, W. Bernstein, Phys. Rev. 98, 1828, 1955; 103, 1253, 1956; [Abstractor's note: This is given as a single reference]; W.P. Jesse, J. Sadauskis,

Card 4/5

27316

P/046/60/005/011/005/018  
D249/D303

Ionization of molecular ...

Phys. Rev., 97, 1668, 1955; *ibid*, 107, 766, 1957; W.L. Fite, L.T. Brackmann, Phys. Rev., 112, 1141, 1958.

ASSOCIATION: Institute of Nuclear Research, Warsaw, Laboratory of Radiation Chemistry; Institute of Physical Chemistry, Warsaw, Department of Electrochemistry

SUBMITTED: September, 1960

X

Card 5/5

KOLOS, W.

On wave functions for the problem of electron and X-ray scattering by  
helium atoms. Bul Ac Pol mat 8 no.1:67-70 '60. (EEAI 9:11)

1. Laboratory of Radiation Chemistry, Institute for Nuclear Research  
Polish Academy of Sciences. Presented by L. Infeld.

(Helium)

(Electrons)

(X rays)

(Eigenfunctions)

21305

P/045/61/020/002/002/006  
B108/B209

24. 4500

AUTHORS:

Kozlos, W., Wolniewicz

TITLE:

Coupling between electronic and nuclear motions, and relativistic effects in the ground state of the  $H_2$  molecule

PERIODICAL:

Acta Physica Polonica, v. 20, no. 2, 1961, 129-140

TEXT: The authors calculated the contributions made by the coupling between nuclear and electronic motion and relativistic effects to the binding energy of the hydrogen molecule, since these contributions must be known for comparing experimental values with the lowest eigenvalue of the nonrelativistic Hamiltonian calculated for the  $H_2$  molecule having a fixed nucleus with the experiment. The coupling between nuclear and electronic motion changes the molecular energy in first approximation by the expression

$$E' = E - E_0 = \int \Psi^* H' \Psi d\tau_1 d\tau_2. \quad (1)$$

(Ref. 2: Dalgarno, A. and McCarroll, R., Proc. Roy. Soc., 237, 383 (1956)).  
The wave function is assumed to be real and given by the expansion

Card 1/6

21305

Coupling between ...

P/045/61/020/002/002/006  
B108/B209

$$\Psi = \sum_i c_i(R) \Psi_i(\xi_i, \eta_i, \xi_2, \eta_2, \varphi_1, \varphi_2), \quad (3)$$

where  $\xi_i$  and  $\eta_i$  are elliptic coordinates with respect to R, and  $\varphi_i$  is the azimuthal angle of the i-th electron. For the calculation of the expectation value of  $\Delta_R$  ( $\vec{R}$  denotes the relative coordinate of the nuclei), this operator is expressed in terms of  $\xi_i, \eta_i, \varphi_i, R, \theta$ , and  $\phi$  ( $\theta$  and  $\phi$  are the polar angles of R). The resulting expression may be simplified if one restricts the problem to wave functions of the form

$$\Psi = \sum c_k^p(R) g_k(\xi_1, \eta_1, \xi_2, \eta_2) \chi^p(\xi_1, \eta_1, \xi_2, \eta_2, \varphi) \quad (5)$$

where  $\varphi = \varphi_1 - \varphi_2$ , and  $R^p \chi^p$  is a homogeneous function of p-th order of the position vector  $\vec{r}_i$  of the i-th electron only. Thus, one obtains

$$\int \Psi \Delta_R \Psi d\tau_1 d\tau_2 = \sum_p \int \Psi R^p \chi^p \Delta_R R^{-p} c_k^p g_k d\tau_1 d\tau_2. \quad (7)$$

where all terms containing derivatives with respect to any angle and, due to the symmetry  $\chi^p(\varphi + 2\pi) = \chi^p(\varphi)$ , all terms linear in  $\cos \varphi_1$  or  $\sin \varphi_1$  may be

Card 2/8

Coupling between ...

21305  
P/045/61/020/002/002/006  
B108/B209

omitted. It is pointed out that for solving Eq. (7) one must know the derivatives  $\frac{d\psi}{dR} = \sum \frac{dc_k}{dR} \psi_k$ . For the calculation of the relativistic correction it is necessary to use the Hamiltonian given by Bethe, H. A. and Salpeter, E. E. (Ref. 1: Encyclopedia of Physics, 35/1, 267 (1957)). Thus,  $E_{\text{rel}} = \int \psi H_{\text{rel}} \psi d\tau = E_1 + E_2 + E_3 + E_4 + E_5 + E_6$ . For the ground state of the  $H_2$  molecule,  $E_3$  and  $E_6$  are zero, and if use is made of the symmetry of  $\psi$ , the non-vanishing contributions to  $E_{\text{rel}}$  read

$$E_1 = -\frac{\hbar^4}{4m^3 c^2} \int (\Delta_n \psi)^2 d\tau_1 d\tau_2,$$

$$E_2 = -\frac{e^2}{2(mc)^2} \int \psi \frac{1}{r_{12}} \left[ p_1 \cdot p_2 + \frac{r_{12}(r_{12} \cdot p_1) p_2}{r_{12}^2} \right] \psi d\tau_1 d\tau_2,$$

$$E_4 = \frac{4\pi^2 e^2 \hbar^2}{(2mc)^2} \int \psi \psi [2\delta(r_{12}) - \delta(r_{13})] d\tau_1 d\tau_2,$$

$$E_5 = \frac{8\pi e^2 \hbar^2}{(2mc)^2} \int \psi \psi \delta(r_{12}) d\tau_1 d\tau_2,$$

(16)

Card 3/6



Coupling between ...

21305  
P/045/61/020/002/002/006  
B108/B209

Results: The wave function the authors employed was of the form

$$\Psi = \frac{1}{2\pi} e^{-\alpha(r_1+r_2)} \sum_{i=1}^n c_i(R) g_i, \quad n = 1, 2, 3, 4, 5, \quad (18)$$

where

$$\begin{aligned} g_1 &= 2, \quad g_2 = \eta_1^2 + \eta_2^2, \quad g_3 = 2\eta_1\eta_2, \\ g_4 &= \xi_1 + \xi_2, \quad g_5 = 2[(\xi_1^2 - 1)(1 - \eta_1^2)(\xi_2^2 - 1)(1 - \eta_2^2)]^{1/2} \cos(\varphi_1 - \varphi_2). \end{aligned} \quad (18)$$

All calculations have been carried out for the internuclear distance  $R = 1.4$  au and using  $\alpha = 0.95$ . The coefficients  $c_i(R)$  as well as  $E_0$ , i. e., the expectation value of  $H_0$ , and the corresponding binding energies  $D$  are given in Table I. The coupling corrections computed with the aid of four wave functions are presented in Table II, where

Card 4/8

Coupling between ...

21305

P/045/61/020/002/002/006  
B108/B209

$$E_1' = -\frac{\hbar^2}{M} \int \Psi \Delta_R \Psi d\tau_1 d\tau_2$$

$$E_2' = -\frac{\hbar^2}{4M} \int \Psi (\Delta_{r_1} + \Delta_{r_2}) \Psi d\tau_1 d\tau_2$$

$$E_3' = -\frac{\hbar^2}{2M} \int \Psi \nabla_{r_1} \nabla_{r_2} \Psi d\tau_1 d\tau_2$$

in au/M (= 119.47 cm<sup>-1</sup>) and E' = E<sub>1</sub>' + E<sub>2</sub>' + E<sub>3</sub>' in cm<sup>-1</sup>. The relativistic corrections have been calculated with the aid of all five wave functions. The results in cm<sup>-1</sup> ( $\frac{a^2}{2}$  au = 5.844 cm<sup>-1</sup>) are presented in Table III, where E'' = E<sub>rel</sub> - E<sub>2</sub>. The results obtained by J. Ladik (Ref. 6: Acta phys. Hungar., 10, 271 (1959)) with the Wang wave function are also given in Table III. Addition of the calculated corrections to the accurate non-relativistic result for infinitely heavy nuclei gives the theoretical binding energy D = 38,280 cm<sup>-1</sup>, the experimental value being D = 38,286 ± 6 cm<sup>-1</sup>. There are 3 tables and 8 non-Soviet-bloc references.

Card 5/6

21305

P/045/61/020/002/002/006  
B108/B209

Coupling between ...

ASSOCIATION: Department of Electrochemistry, Institute of Physical  
Chemistry, Polish Academy of Sciences, Warsaw and Department  
of Theoretical Physics, N. Copernicus University Toruń

SUBMITTED: July 21, 1961

n	$E_g$ (au)	D (eV)	$c_1$	$c_2$	$c_3$	$c_4$	$c_5$
1	-1.08612	2.343	4.07284				
2	-1.11544	3.141	3.45343	2.03585			
3	-1.13560	3.690	3.38893	2.16629	-1.47897		
4	-1.15165	4.126	1.73628	2.00821	-1.38824	0.82506	
5	-1.16081	4.376	1.70621	1.95257	-1.35017	0.84280	-0.16518

Table 1

Card 6/6

21308

P/045/61/020/002/006/006  
B108/B209

26.2212

AUTHOR: Kołos, Włodzimierz

TITLE: Approximate wave functions in the calculation of electron excitation accompanying the beta decay of  $\text{He}^6$

PERIODICAL: Acta Physica Polonica, v. 20, no. 2, 1961, 175-178

TEXT: The purpose of the present study is to show the probability of non-excitation of atomic electrons in the beta-decay of  $\text{He}^6$ . In publications the suggestion has been made to use self-consistent field wave functions in the calculation of the mean beta-induced ionization in order to improve the results. The main effects in beta decay come from the shaking of the nucleus due to a change in charge. In perturbation theory, the probability of finding the daughter  $\text{Li}^+$  ion in its n-th quantum state is, given by  $P_{0n} =$

$\left| \int \psi_0 \psi_n^* d\tau \right|^2$ , where  $\psi_0$  denotes the ground-state wave function of the helium atom, and  $\psi_n$  the wave function for the n-th state of the  $\text{Li}^+$  ion.  $\sum_n P_{0n} = 1$ , where  $\sum_n$  indicates summation over all states in the discrete spectrum or

Card 1/5

21308

P/045/61/020/002/006/006  
B108/B209

Approximate wave ...

integration over the continuous spectrum. The author calculated  $P_{00}$  (for the ground state) for He with the following wave functions obtained by the self-consistent field method:

$$\varphi_{01} = N_1 e^{-\zeta(r_1+r_2)} \quad (3)$$

$$\varphi_{02} = N_2 \chi(r_1) \chi(r_2), \quad (4)$$

$$\varphi_{03} = N_3 [e^{-\zeta_1 r_1 - \zeta_2 r_2} + e^{-\zeta_2 r_1 - \zeta_1 r_2}], \quad (5)$$

$$\varphi_{04} = N_4 e^{-\zeta(r_1+r_2)} (1 + c_{12}), \quad (6)$$

$$\varphi_{05} = N_5 e^{-\zeta(r_1+r_2)} [1 + c_1 r_{12} + c_2 (r_1 - r_2)^2], \quad (7)$$

where  $\chi$  represents the self-consistent field orbitals computed by Weiss, A. (Ref. 6: Unpublished results, The University of Chicago (1958)) in the form of expansion in terms of the Laguerre polynomials

$$\chi(r) = e^{-\zeta r} \sum_{n=1}^{\infty} a_n L_{n+1}^1(2\zeta r). \quad (8)$$

For the  $\text{Li}^+$  ion, the functions  $\psi_{0n}$ , which have the same form as  $\varphi_{0n}$ , have

Card 2/6

21308

P/045/61/020/002/006/006  
B108/B209

Approximate wave ...

been employed. In Eq. (8), a five-term expansion and, for helium, also a three-term expansion (corresponding wave function  $\psi_{02}$ ) have been used. The six- and seven-term wave functions used by Winther, A. (Ref. 7: K. Danske Vidensk. Selsk. Mat.-fys. Medd., 27, No. 2 (1952)) for He and  $\text{Li}^+$  are denoted by  $\psi_{0w}$  and  $\psi_{0w}$ , respectively. The results achieved for  $P_{00}$  and for the total energy of both systems are given in Table 1. As the  $P_{00}$  results are very unsatisfactory, the author computed the expectation values of  $r$  and  $r^2$  for helium, using the wave function  $\psi_{01}, \dots, \psi_{05}$  (Table 2).  $r_{ac}$  denotes the accurate expectation value. In this case, the self-consistent field method yields good results. The author points out that the probability  $P_{00}$  should be calculated with much more accurate wave functions before drawing conclusions on the applicability of the self-consistent field wave method to this problem. The author thanks A. Weiss for sending him the unpublished results of his self-consistent field calculations. There are 4 tables and 7 non-Soviet-bloc references.

Card 3/4

APPROVED FOR RELEASE: 09/18/2001

CIA-RDP86-00513R000823920014-

P/045/61/020/002/006/006  
B108/B209

Approximate wave ...

ASSOCIATION: Laboratory of Radiation Chemistry, Institute of Nuclear Research, Polish Academy of Sciences, Warsaw and Department of Electrochemistry, Institute of Physical Chemistry, Polish Academy of Sciences, Warsaw

SUBMITTED: September 15, 1960

Table I

Values of  $P_{00}$  calculated with various wave functions

Wave functions	$\psi_{01}, \psi_{01}$	$\psi_{02}, \psi_{02}$	$\psi_{03}, \psi_{03}$	$\psi_{04}, \psi_{04}$	$\psi_{05}, \psi_{05}$	$\psi_{06}, \psi_{06}$	$\psi_{07}, \psi_{07}$
$P_{00}$	0.7247	0.7317	0.7314	0.7110	0.7174	0.6859	0.6698
— Energy (au)	10.0700	10.0980	10.0981	10.1247	10.1516	10.1707	10.1827

Card 4/4

KOLOS, Wlodzimierz; KOSEK, Stanislaw

Cerenkov radiation in the  $^{60}\text{Co}$  gamma irradiation unit.  
Mikleonika 7 no.6:379-388 '62.

1. Institute of Nuclear Research, Polish Academy of Sciences,  
Warsaw, Department of Radiation Chemistry.

KOLOS, Włodzimierz

On the various forms of hydrogen atoms appearing in the gamma irradiated aqueous solutions. Nukleonika 8 no.7:451-463 '63.

1. Department of Radiation Chemistry, Institute of Nuclear Research, Warsaw.